Evolve 480 SC

Reg. No. L10644 (Act 36 of 1947) Warning. Harmful if swallowed, May be harmful in contact with skin, Very toxic to aquatic life with long lasting effects.

Always adhere to label



12 - 24 Curative

Protective

Composition:

- Azoxystrobin 240 g/ℓ (FRAC 11) + Fludioxonil, 240 g/ℓ (FRAC 12)

Recommended Application & Dosage:

- Pre-packhouse drench / bath or in-line drench: 250 mℓ/100 ℓ water for 600 ppm azoxystrobin & 600 ppm fludioxonil
- Wax application: 375 mℓ in 25 ℓ wax at ~ 1 ℓ of wax mixture per 1 ton of fruit

Top Up:

- Apply at a rate of 250 mℓ Evolve 480 SC for every 100 ℓ of fresh water
- Top up with 7 mℓ Evolve 480 SC to water mixture for every ton of fruit treated

Purpose:

- To control blue & green mould (*Penicillium* spp.)
- For effective *Penicillium* sporulation suppression
- To control imazalil and/or pyrimethanil resistant green mould

Available in:

1 ℓ

Effective residue loading:

Azoxystrobin	Fludioxonil
0.4 – 1.5 mg/kg	0.8 – 1.5 mg/kg

- Depends on fruit type and application

Tips:

- Keep mixture clean
- Constant agitation of bath required
- Can be applied in a heated mixture