ICA Thiabendazole 500 SC

Reg. No. L7784 (Act 36 of 1947) Warning. May be harmful if swallowed, May be harmful in contact with skin, Very toxic to aquatic life with long lasting effects.

Always adhere to label



24 h Curative

Protective

Composition:

- Thiabendazole 500 g/ℓ (FRAC 1)

Recommended Application & Dosage:

- Pre-packhouse drench / bath or in-line drench: 200 mℓ/100 ℓ water for 1000 ppm thiabendazole
- Wax: 200 mℓ/25 ℓ wax for 4000 ppm thiabendazole

Top Up:

- Apply at a rate of 200 mℓ ICA Thiabendazole 500 SC for every 100 ℓ of fresh water added
- Top up with 5 mℓ of ICA Thiabendazole 500 SC to water mixture for every ton of fruit treated

Purpose:

- To control blue & green mould (*Penicillium* spp.)
- To control latent pathogens stem-end rot

Available in:

5 ℓ

Effective residue loading:

1 mg/kg

- Depends on fruit type and application

Tips:

- Keep mixture clean
- Wax and water mixtures must be continuously agitated
- Often applied in pre-packhouse drench and wax
- Reduction of chilling injury on cold stored fruit according to literature
- Two water applications rates available, for double and single application