ImazaCure 750 SG

Reg. No. L7926 (Act 36 of 1947) Warning. Harmful if swallowed, May cause an allergic skin reaction, Very toxic to aquatic life with long lasting effects.

Always adhere to label



24 h Curative

Protective

Composition:

- Imazalil as Sulphate 750 g/kg (FRAC 3)

Recommended Application & Dosage:

- Pre-packhouse drench / bath or in-line drench: 670 g/1 000 ℓ water for 500 ppm Imazalil

Top Up:

- Apply at a rate of 67 g ImazaCure 750 SG for every 100 ℓ fresh water
- Top up with 5 g ImazaCure 750 SG to water mixture for every ton of fruit treated

Purpose:

- To control blue & green mould (*Penicillium* spp.) decay and sporulation suppression

Tips:

- Keep mixture clean
- Adjust the water pH to ≤ 7 before adding ImazaCure 750 SG (only where water quality is poor)
- Final mixture pH must be < 6,5
- When used in combination with EcoTizer, pH must be managed between 5,5 - 6,5.
- The EcoTizer test kit and Imazacure titration kit can be used in mixtures where both are present.

Available in:

670 g

Effective residue loading:

2 mg/kg

- Depends on fruit type and application

Test Kit Available:



WATCH INSTRUCTION VIDEO