StrobiCure 250 SC

Reg. No. L10557 (Act 36 of 1947) Warning. Harmful if inhaled, May cause an allergic skin reaction, Very toxic to aquatic life with long lasting effects.

Always adhere to label



12 h Curative

Protective

Composition:

- Azoxystrobin 250 g/ℓ (FRAC 11)

Recommended Application & Dosage:

- Pre-packhouse drench / bath or in-line drench: 450 mℓ/100 ℓ water for 1125 ppm azoxystrobin
- Wax application: 400 mℓ in 25 ℓ wax at ~ 1 ℓ of wax mixture per 1 ton of fruit

Top Up:

- Apply at a rate of 450 mℓ StrobiCure 250 SC for every 100 ℓ of fresh water
- Top up with 10 mℓ StrobiCure 250 SC to water mixture for every ton of fruit treated

Purpose:

- To control postharvest decay caused by green mould (*Penicillium digitatum*) and Phytophthora brown rot (*Phytophthora nicotianae*)
- Postharvest sporulation control of green mould (wax only application, control of ≥ 50% expected)

Available in:

5 ℓ

Effective residue loading:

0.4 – 1.5 mg/kg

- Depends on sampling point, fruit type and application

Tips:

- Use in combination with fungicide from another FRAC group to minimize resistance risk
- Agitate treatment mixture continuously
- Can be applied in a heated mixture